

学校编码: 10384

分类号_____密级_____

学 号: 19220051301772

UDC_____

厦 门 大 学

硕 士 学 位 论 文

芳樟叶黄酮和多糖的提取分离、结构鉴定及
抗氧化活性研究

Extraction, Separation, Identification and Antioxidant
activities of Flavonoid and Polysaccharide from
Cinnamomum Camphora Leaves

王 先

指导教师姓名: 李 清 彪 教 授

王 远 鹏 助 理 教 授

专 业 名 称 : 生 物 化 工

论文提交日期: 2 0 0 8 年 9 月

论文答辩日期: 2 0 0 8 年 9 月

学位授予日期: 2 0 0 8 年 月

答辩委员会主席: _____

评 阅 人: _____

2008 年 8 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

- () 1. 经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。
- () 2. 不保密，适用上述授权。

声明人（签名）：

年 月 日

摘要

我国是世界上生产芳樟油最多的国家，其产量占世界的80%。樟树（*Cinnamomum camphora*）是我国生产天然香料芳香油的主要树种，经济价值高。近年来，通过组培、扦插等方法培育出纯种芳樟，大大提高樟树资源的经济价值。但在提取精油过程中产生数量巨大的芳樟枝叶等废弃物，由于对芳樟废弃物的资源化研究力度不够，忽视了许多有药用价值的化学成分如黄酮、多糖等化合物的开发利用，造成芳樟资源浪费严重，有必要改进和发展芳樟中有效成分的提取纯化工艺以及寻找其中有生物活性的化学成分，为芳樟资源的综合利用提供理论与实践依据，发挥其最好的社会效益和经济效益。

通过对芳樟树不同部位的测定发现，不同部位中总黄酮含量和FRAP值均有明显差异，其中芳樟叶的总黄酮含量和FRAP值最高。不同部位中总黄酮含量与FRAP值呈正相关性。高效液相色谱法（HPLC）测定结果表明芳樟叶中黄酮苷元主要含有槲皮素、山奈素和异鼠李素，其含量分别为：0.39%、0.14%和0.063%，在芳樟叶中大部分以黄酮糖苷的形式存在。

通过单因素实验和正交试验优化了芳樟叶黄酮的微波辅助萃取工艺条件：乙醇浓度60%（v/v），料液比1:12（g/mL），微波功率320 W，间歇辐射两次，每次1 min，芳樟叶黄酮提取率达2.97%，比乙醇热回流提取提高了6.83%，并且提取时间缩短了99%；为了进一步纯化芳樟叶黄酮，通过测定10种大孔树脂对芳樟叶黄酮的静态吸附量和解吸率，确定了芳樟叶黄酮纯化工艺的吸附剂DA201树脂，并优化了DA201树脂对芳樟叶黄酮的动态吸附和解吸工艺条件：上柱液黄酮浓度为13.20 mg/mL，树脂柱径高比为1:10，吸附流速为3 BV/h；洗脱溶剂为5 BV 50%的乙醇水溶液，洗脱流速为3 BV/h。芳樟叶粗提液经装填有大孔树脂DA201的固定床纯化后黄酮含量由38.15%提高到91.18%，纯化倍数2.39倍。

采用HPLC法测定芳樟叶精制黄酮中芦丁、槲皮素、山奈酚和异鼠李素的含量，分别为123.33 mg/g、70.00 mg/g、28.63 mg/g和20.81 mg/g。利用高效液相色谱—二极管阵列检测器—电喷雾质谱（HPLC-DAD-ESI-MS）联用技术对芳樟叶精制黄酮的主要组分进行了定性分析，初步鉴定出芦丁、槲皮素-3-O- β -D-半乳糖苷、山奈酚-3-O- β -D-芸香糖苷、异鼠李素-3-O- β -D-芸香糖苷、槲皮素-3-O- β -D-鼠李糖苷、山奈酚-3-O- β -D-鼠李糖苷6种黄酮糖苷。采用Sephadex LH-20

凝胶柱从精制黄酮中分离得到芦丁和山奈酚-3-O- β -D-芸香糖苷。

进一步采用沸水提取、乙醇沉淀的方法，从芳樟叶中提取得到水溶性粗多糖 CLPS。经DEAE52-纤维素柱和Sephacryl-S400凝胶柱分离得到一种酸性多糖组分 CLPS-A2，采用高效凝胶渗透色谱法测定其为均一性多糖，平均分子量为244 KDa。紫外可见吸收光谱和红外光谱的分析结果表明CLPS-A2为具有乙酰氨基结构的 β 型吡喃多糖。CLPS-A2水解产物的分析结果显示，CLPS-A2是一种主要由鼠李糖、木糖、阿拉伯糖、葡萄糖、半乳糖组成的酸性杂多糖。

利用 FRAP 法和 DPPH 法测定了芳樟叶各提取物的体外抗氧化活性。其中，芳樟叶总黄酮苷元的抗氧化活性最强，其次是芳樟叶精制黄酮，芳樟叶多糖的抗氧化活性最弱。结合对黄酮结构的分析可知，芳樟叶的抗氧化活性与其所含的黄酮醇苷有关。

关键词：芳樟叶；黄酮；多糖；提取；分离；结构鉴定；抗氧化活性

ABSTRACT

Cinnamomum camphora is an important plant that has been used to produce natural flavor and aromatic oils in China for a long time. In recent years, it was reported that purebred *Cinnamomum camphora* could be successfully cultivated by tissue culture, which significantly enhanced the value of *Cinnamomum camphora*. However, considerable components of pharmacological merit in *Cinnamomum camphora*, e.g. flavonoid, polysaccharide, received little attention. To utilize *Cinnamomum camphora* trees more reasonably and economically, the techniques for extracting pharmacological components from *Cinnamomum camphora* trees should be improved and exploited.

A large variation of total flavonoid contents and FRAP values in different parts of *Cinnamomum camphora* trees were observed. The *Cinnamomum camphora* leaves showed to be of the highest total flavonoid content and FRAP value. A positive linear correlation between FRAP values and total flavonoid contents were established. After the acid-catalyzed hydrolysis of the leaves quercetin(0.39%), kaempferol(0.14%) and isorhmnetin(0.063%) were detected and founded to exist mainly in the form of flavonoid glucosides.

By a microwave-assisted extraction technology flavonoid was extracted from the residue of *Cinnamomum camphora* leaves after essential oil extraction. Macroporous resin adsorption was used for further purification. With single factor and orthogonal designed experiments the optimal extraction conditions were determined as follows: 60% ethanol aqueous solution, ratio of solid to liquid 1:12(g/mL), microwave power 320 W and treated for 2 min,. Under such conditions, the yield of the flavonoid reached 2.97%, which was 6.82% higher than that obtained using heat ethanol reflux extraction, and the extraction operation duration of the technology was significantly reduced to 1% of that for the control process. Ten kinds of macroporous resin were then examined for further purification of *Cinnamomum camphora* flavonoid. DA201 resin was showed to possess the best absorption and desorption behavior. The optimum adsorption and desorption conditions were sample solution flavonoid

concentration 13.20 mg/mL, diameter to height 1:10, absorption rate 3BV/h; The optimum desorption conditions were volume of 50% ethanol 150 mL (approximately 5 BV) as eluting solvent, eluting rate 3BV/h. The content of flavonoid could be improved to 91.18% from 38.15% after an absorption-desorption operation applied upon a DA201 resin packed column.

By HPLC, rutin(123.33 mg/g), quercetin(70.00 mg/g), kaempferol(28.63 mg/g), isorhamnetin(20.81 mg/g) were identified and determined in the purified flavonoid obtained above. By HPLC-DAD-ESI-MS, six flavonoid glycosides, rutin, quercetin-3-O- β -D-galactoside, kaempferol-3-O- β -D-rutinoside, isorhamnetin-3-O- β -D-rutinoside, quercetin-3-O-rhamnoside and kaempferol-3-O- β -D-rhamnoside were primarily identified in the purified flavonoid. Rutin and kaempferol-3-O- β -D-rutinoside were separated from the purified flavonoid by sephadex LH-20 column chromatography.

The crude water-soluble polysaccharide (CLPS) was obtained from *Cinnamomum camphora* leaves by boiled water extraction and ethanol precipitation. DEAE52-cellulose column and Sephacryl-S400 gel filtration column were used to further separate an acid polysaccharide (CLPS-A2) from the crude polysaccharide. The molecular weight of CLPS-A2 was determined to be 244 kDa. UV-VIS spectrum and FTIR analysis showed that CLPS-A2 was β -heterosaccharides with pyran and acetyl amino group. Analysis of the hydrolysis products of CLPS-A2 showed that CLPS-A2 was a heterosaccharide composed of rhamnose, arabinose, xylose, glucose and galactose.

The antioxidant activities of various extracts from *Cinnamomum camphora* leaves were evaluated by FRAP assay and DPPH assay in vitro. The result showed that the antioxidant activities of various extracts ranged as the following order: total flavonoid-aglycone>purified flavonoid>polysaccharide. It was inferred that the potent antioxidant activities of *Cinnamomum camphora* leaves were mainly attributed to flavonol glycosides.

Key words: *Cinnamomum camphora*; flavonoid; polysaccharide; extraction; separation; identification; antioxidant activities

目录

第一章 文献综述	1
1.1 樟树简介	1
1.2 樟树化学成分及药理作用	2
1.2.1 樟树的化学成分	2
1.2.2 樟树的药理作用	4
1.3 黄酮类化合物的研究进展	6
1.3.1 黄酮类化合物的概述	6
1.3.2 黄酮类化合物的提取分离	9
1.3.3 黄酮类化合物的分析检测方法	13
1.3.4 黄酮类化合物的生物活性	14
1.4 多糖的研究进展	16
1.4.1 多糖的概述	16
1.4.2 多糖的提取分离	16
1.4.3 多糖的纯度与分子量检测	19
1.4.4 多糖的结构分析	20
1.5 本文的研究目的、意义及内容	21
第二章 芳樟树不同部位中总黄酮含量及其抗氧化活性	23
2.1 引言	23
2.2 材料与方法	23
2.2.1 试剂及仪器	23
2.2.2 实验材料	23
2.2.3 提取过程	23
2.2.4 测定方法	23
2.3 结果与讨论	24
2.3.1 标准曲线	24
2.3.2 芳樟树不同部位中总黄酮含量及其抗氧化活性	25
2.3.3 芳樟叶中黄酮苷元的含量	26

2.4 本章小结	27
第三章 芳樟叶黄酮的提取与纯化	28
3.1 引言	28
3.2 材料与方法	28
3.2.1 试剂及仪器	28
3.2.2 实验材料	28
3.2.3 芳樟叶主要成分分析	28
3.2.4 芳樟叶黄酮的提取工艺	29
3.2.5 芳樟叶黄酮的纯化工艺	30
3.3 结果与讨论	31
3.3.1 芳樟叶的主要成分	31
3.3.2 微波辅助萃取工艺的优化	32
3.3.3 大孔树脂纯化工艺的优化	36
3.4 本章小结	42
第四章 芳樟叶黄酮的分离与结构鉴定	43
4.1 引言	43
4.2 材料与方法	43
4.2.1 试剂及仪器	43
4.2.2 实验材料	43
4.2.3 HPLC检测	43
4.2.4 HPLC-DAD-ESI-MS分析	44
4.2.5 Sephadex LH-20 柱分离	44
4.2.6 单体化合物的结构鉴定	45
4.3 结果与讨论	45
4.3.1 精制黄酮中芦丁的含量	45
4.3.2 精制黄酮中黄酮苷元的含量	46
4.3.3 精制黄酮中主要组分的定性分析	46
4.3.4 Sephadex LH-20 柱对精制黄酮的分离效果	48
4.3.5 单体化合物的结构分析	50

4.4 本章小结	53
第五章 芳樟叶多糖的提取、分离与结构鉴定	54
5.1 引言	54
5.2 材料与方法	54
5.2.1 试剂及仪器	54
5.2.2 实验材料	54
5.2.3 多糖的提取	54
5.2.4 多糖的分离	54
5.2.5 多糖纯度鉴定及分子量测定	55
5.2.6 多糖的结构鉴定	57
5.3 结果与讨论	57
5.3.1 粗多糖的理化性质	57
5.3.2 DEAE52-纤维素柱对粗多糖分离效果	58
5.3.3 Sephacryl-S400 凝胶柱对多糖CLPS-A的分离效果	59
5.3.4 多糖CLPS-A2 的结构分析	60
5.4 本章小结	63
第六章 芳樟叶提取物体外抗氧化活性	64
6.1 引言	64
6.2 材料与方法	64
6.2.1 试剂及仪器	64
6.2.2 实验材料	64
6.2.3 芳樟叶提取物的制备	64
6.2.4 FRAP法	65
6.2.5 DPPH法	65
6.3 结果与讨论	66
6.3.1 总抗氧化能力	66
6.3.2 清除DPPH自由基能力	67
6.4 本章小结	68
结论与建议	70

参考文献.....	72
附录.....	80
在学期间所发表论文.....	86
致谢.....	87

厦门大学博硕士论文摘要库

Contents

Chapter 1 Introduction	1
1.1 Introduction of <i>Cinnamomum camphora</i>	1
1.2 Chemical constituents and pharmacological activities of <i>Cinnamomum camphora</i>	2
1.2.1 Chemical constituents of <i>Cinnamomum camphora</i>	2
1.2.2 Pharmacological activities of <i>Cinnamomum camphora</i>	4
1.3 Research progress on flavonoid.....	6
1.3.1 Introduction of flavonoid.....	6
1.3.2 Extraction and separation of flavonoid	9
1.3.3 Quantification and identification of flavonoid	13
1.3.4 Pharmacological activities of flavonoid	14
1.4 Research progress on polysaccharide.....	16
1.4.1 Introduction of polysaccharide.....	16
1.4.2 Extraction and separation of polysaccharide.....	16
1.4.3 Polysaccharide purity and molecular weight determination	19
1.4.4 Identification of polysaccharide	20
1.5 Goal, significance and content of this study.....	21
Chapter 2 Determination of total flavonoids and their antioxidant activities in different parts of <i>Cinnamomum camphora</i>	23
2.1 Introduction	23
2.2 Material and methods	23
2.2.1 Reagents and equipments	23
2.2.2 Experiment materials.....	23
2.2.3 Extracting process	23
2.2.4 Determination methods	23
2.3 Results and discussion.....	24

2.3.1 Standard curves	24
2.3.2 Total flavonoid contents and antioxidant activities in different parts of <i>Cinnamomum camphora</i>	25
2.3.3 Flavonoid aglycone content in <i>Cinnamomum camphora</i> leaves.....	26
2.4 Conclusions	27
Chapter 3 Extraction and purification of flavonoid from <i>Cinnamomum camphora</i> leaves	28
3.1 Introduction	28
3.2 Material and methods	28
3.2.1 Reagents and equipments	28
3.2.2 Experiment materials.....	28
3.2.3 Analysis on main components of <i>Cinnamomum camphora</i> leaves.....	28
3.2.4 Technology of extracting flavonoid from <i>Cinnamomum camphora</i> leaves.....	29
3.2.5 Technology of purifying flavonoid from <i>Cinnamomum camphora</i> leaves.....	30
3.3 Results and discussion.....	31
3.3.1 main components of <i>Cinnamomum camphora</i> leaves.....	31
3.3.2 Optimization of processing parameters for micro-wave assisted extracion.....	32
3.3.3 Optimization of processing parameters for purification by macroporous resin.....	36
3.4 Conclusions	42
Chapter 4 Separation and structure identification of flavonoid glycosides from <i>Cinnamomum camphora</i> leaves.....	43
4.1 Introduction	43
4.2 Material and methods	43
4.2.1 Reagents and equipments	43
4.2.2 Experiment materials.....	43
4.2.3 Determination by HPLC.....	43
4.2.4 Analysis by HPLC-DAD-ESI-MS	44
4.2.5 Separation on Sephadex LH-20.....	44

4.2.6 Structure identification of monomeric compound.....	45
4.3 Results and discussion	45
4.3.1 Content of rutin in purified flavonoid	45
4.3.2 Content of flavonoid aglycone in purified flavonoid	46
4.3.3 Structure analysis of main components in purified flavonoid	46
4.3.4 Effect of Sephadex LH-20 on separation of purified flavonoid	48
4.3.5 Structure analysis of monomeric compound	50
4.4 Conclusions	53
Chapter 5 Extraction, separation and structure identification	
of polysaccharide from <i>Cinnamomum camphora</i> leaves	54
5.1 Introduction	54
5.2 Material and methods	54
5.2.1 Reagents and equipments	54
5.2.2 Experiment materials.....	54
5.2.3 Extraction of polysaccharide.....	54
5.2.4 Separation of polysaccharide.....	54
5.2.5 Determination of polysaccharide purity and molecular Weight	55
5.2.6 Structure identification of polysaccharide.....	57
5.3 Results and discussion	57
5.3.1 Physicochemical property of crude polysaccharide	57
5.3.2 Effect of DEAE 52-cellulose column on separation of crude polysaccharide ...	58
5.3.3 Effect of Sephacryl-S400 gel column on separation of polysaccharide CLPS-A	
.....	59
5.3.4 Structure analysis of polysaccharide CLPS-A2	60
5.4 Conclusions	63
Chapter 6 Antioxidant activities of various extracts from	
<i>Cinnamomum camphora</i> leaves in vitro	64
6.1 Introduction	64

6.2 Material and methods	64
6.2.1 Reagents and equipments	64
6.2.2 Experiment materials	64
6.2.3 Preparation of various extracts from <i>Cinnamomum camphora</i> leaves	64
6.2.4 FRAP assay	65
6.2.5 DPPH assay	65
6.3 Results and discussion	66
6.3.1 Total antioxidant activity	66
6.3.2 Scavenging DPPH free radical ability	67
6.4 Conclusions	68
Conclusions and suggestions	70
References	72
Appendix	80
Publications	86
Acknowledgement	87

第一章 文献综述

1.1 樟树简介

樟树 (*Cinnamomum camphora*), 别名: 香樟、樟木、乌樟、油樟、香樟、芳樟。樟树是樟科 (*Lauraceae*) 樟属 (*Cinnamomum*) 植物, 系常绿乔木, 单株高度可达 30 米, 树冠广卵形; 树皮幼时绿色, 平滑, 老时渐变为黄褐色或灰褐色纵裂, 冬芽卵圆形。叶互生, 革质薄, 呈卵状椭圆形, 长 6~12 厘米, 宽 3~6 厘米, 顶端短尖或近尾尖, 基部圆形, 背面呈灰绿色, 两面无毛, 有离基三出脉, 近叶基的第一对或第二队侧脉长而显著, 脉腋有明显的腺体。圆锥花序腋生于新枝, 长 5~7.5 cm; 花小, 淡黄绿色; 花被 6 片, 椭圆形, 长约 2 毫米, 内面密生短柔毛; 能育 9 个雄蕊, 花药 4 室; 第三轮雄蕊花药外向瓣裂; 子房球形, 无毛。核果球形, 直径 6-8 mm, 熟时紫黑色; 果托杯状。花期 5 月, 果实 9-11 月成熟。

樟树是我国生产天然香料芳香油的主要树种, 同时也是珍贵用材树种及园林绿化的优良树种。樟树是热带和亚热带常绿阔叶林的代表树种, 被誉为江南宝树。其分布区广, 分布区域在北纬 10°~34°, 东经 88°~120°, 主产于我国海南、台湾、福建、江西、广东、广西、湖北、湖南、四川、重庆、云南、贵州、浙江等省区, 以福建、台湾为最多。樟树的栽培利用早在东周春秋时代就有记载, 至今已有三千多年的栽培历史。樟树用途广, 经济价值高, 可谓全身都是宝, 其主副产品广泛应用于建筑、工艺、化工、军工、医药等方面。其木材致密美观, 具有芳香味, 抗虫蛀, 可供建筑、造船、家具、乐器、手工艺品之用; 樟树的茎叶均可提制樟脑、樟油, 油中含有桉叶素、黄樟素、芳樟醇、松油醇、柠檬醛等多种重要成分, 是国防、医药、化工香料等工业的重要原料。芳樟油减压精馏所得的第二馏分芳油, 其主要成分是芳樟醇, 为许多高贵香料的主要原料, 既可直接用于调香, 又可制造柠檬醛合成紫罗兰香酮, 还可制天竺葵醇和乙酸芳樟酯等。樟树树姿秀丽、四季常青、枝叶茂密, 具有很强的吸尘能力的抗煤烟能力; 且系深根性树种, 根系特别强大, 主根尤为发达, 有强的抗风能力; 且能吸湿耐水、防风固沙、保护堤岸, 近年来被广泛选作南方城镇及“四旁”绿化树种和环保优良树种。

樟树是樟科植物中枝叶含芳樟醇最高的一种。芳樟醇原料及其派生衍生物的

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库